

NGC 3065: A CERTIFIED LINER WITH BROAD, VARIABLE BALMER LINES

MICHAEL ERACLEOUS¹

Department of Astronomy and Astrophysics, The Pennsylvania State University, 525 Davey Lab, University
 Park, PA 16802

AND

JULES P. HALPERN¹

Department of Astronomy, Columbia University, 550 West 120th St., New York, NY 10027

To Appear in the Astrophysical Journal

ABSTRACT

Motivated by the X-ray properties of the galaxy NGC 3065, we have obtained new optical spectra which reveal that it has a low-ionization nuclear emission-line region (LINER) as well as broad Balmer emission lines, establishing it as an active galactic nucleus. We also examined an older spectrum from the CfA Redshift Survey which, lacking broad Balmer lines, indicates that they appeared some time after 1980. Thus NGC 3065 joins the set of LINERs with broad, variable Balmer lines, which includes such well-known galaxies as NGC 1097 and M81. Inspired by the sometimes double-peaked profiles of the variable Balmer lines in other LINERs, we speculate that the broad Balmer lines of NGC 3065 also come from an accretion disk. We illustrate the plausibility of this hypothesis by fitting a disk model to the observed H α profile. We also estimate the mass of the central black hole as $(9 \pm 4) \times 10^7 M_{\odot}$ from the properties of the host galaxy, which leads to the conclusion that the accretion rate is only $\sim 2 \times 10^{-4}$ times the Eddington value, a property that appears to be common among LINERs. At such a low relative accretion rate the inner accretion disk can turn into a vertically-extended ion torus, which can illuminate the outer, thin disk and power the broad-line emission. The reason for the sudden appearance of broad Balmer lines is an open question, although we suggest two possible explanations: tidal disruption of a star or a sudden transition in the structure of the accretion disk.

Subject headings: galaxies: active—galaxies: individual (NGC 3065)—line profiles

1. INTRODUCTION

Low-ionization nuclear emission-line regions (LINERs; Heckman 1980) are a heterogeneous population of objects, with low-luminosity active galactic nuclei (LLAGNs) and compact starbursts being two of the many possible power sources (see, for example Filippenko 1996, for a review). From the point of view of the physics of accretion onto compact objects, those LINERs that are true LLAGNs are particularly interesting. At the low accretion rates onto the supermassive black holes at the centers of these LINERs, the accretion flows are *qualitatively* different from what occurs in more luminous Seyfert galaxies and quasars. It is therefore extremely interesting that in the past few years several LINERs have been identified as *bona-fide* AGNs based on the presence of broad, and sometimes variable, Balmer lines in their optical spectra (e.g., Ho et al. 1997b). Here we report our discovery of a previously unrecognized LINER and a true AGN with broad and variable Balmer lines in the S0 galaxy NGC 3065 (also known as VII Zw 303). Thus NGC 3065 joins the growing set of LINERs with variable, broad Balmer lines, which includes such famous objects as NGC 1097 (Storch-Bergmann, Baldwin, & Wilson, 1993) and M81 (Bower et al. 1996).

NGC 3065 ($z = 0.00667$, at a distance of 47.3 Mpc; from Tully 1987, but for $H_0 = 50 \text{ km s}^{-1} \text{ Mpc}^{-1}$) has been known to have emission lines since the mid-1950s (Hu-

mason, Mayall, & Sandage 1956; Burbidge & Burbidge 1965). It was detected as a radio source at 1.4 GHz in the NVSS Survey (Condon et al. 1998) with a monochromatic luminosity of $1.2 \times 10^{21} \text{ W Hz}^{-1}$, and as an X-ray source with the *Einstein* IPC with a 0.2–4 keV luminosity of $2.1 \times 10^{41} \text{ erg s}^{-1}$ (Fabbiano, Kim, & Trinchieri 1992). More recent X-ray observations with *ASCA* gave a 2–10 keV luminosity of $5 \times 10^{41} \text{ erg s}^{-1}$, with a spectrum that can be described either as a power-law with a photon index of 1.8, or a 6 keV thermal plasma (Iyomoto et al. 1998). There was no evidence for X-ray emission from a cooler thermal plasma as is found in many other LINERs (e.g., Ptak et al. 1999; Terashima et al. 2000). The hard X-ray spectrum led Iyomoto et al. (1998) to suggest that NGC 3065 harbors a LLAGN and motivated us to obtain new optical spectra in order to evaluate its credentials further.

2. OBSERVATIONS AND SPECTRA: NEW AND OLD

We obtained spectra of NGC 3065 with the MDM Observatory's 2.4m telescope and Boller and Chivens CCD Spectrograph on 2000 May 31 UT, and with the Kitt Peak National Observatory's 2.1m telescope and GoldCam spectrograph on 2000 June 5 UT. In the former set of observations we used a 1''5 slit and a 150 mm⁻¹ grating with a total exposure time of 2000 s to cover the wavelength range 3200–6860 Å at a spectral resolution of 12.4 Å. In the latter we used a 1''9 slit and a 600 mm⁻¹ grating with a total ex-

¹ Visiting astronomer, Kitt Peak National Observatory, which is operated by AURA, Inc., under a cooperative agreement with the National Science Foundation

posure time of 1800 s to cover the wavelength range 5570–8540 Å at a spectral resolution of 4.2 Å. In both cases, spectra were extracted from a $1''.6$ window along the slit. Because of the narrow apertures used, which were comparable to the size of the seeing disk, the absolute flux scale is somewhat uncertain. Wavelength calibration was carried out with the help of arc spectra obtained immediately after the object exposure, and flux calibration was carried out with the help of standard stars observed on the same night and reduced in the same manner as the object. The final, reduced, and combined spectra are shown in Figure 1. To investigate possible variability of the *broad* H α line we also examined the spectrum of NGC 3065 taken during the CfA Redshift Survey on 1980 February 10 with the Mt. Hopkins 1.5m telescope through a $3'' \times 12''$ aperture and originally reported by Tonry & Davis (1981). Although the flux scale of this spectrum was not calibrated, it can still provide information on the presence and profile of any *broad* H α line.




FIG. 1.— Spectra of NGC 3065 obtained at MDM and KPNO in 2000 June. The 3200–8600 Å spectrum was obtained with the MDM 2.4m telescope and has a resolution of 12.4 Å. The lower trace shows the residual spectrum after subtracting the continuum (see §3 of the text for details) with the emission lines identified. The inset shows the H α region of the higher-resolution (4.2 Å) spectrum obtained with the KPNO 2.1m telescope.

3. MEASURED SPECTROSCOPIC PROPERTIES

Although the optical continuum of NGC 3065 is dominated by starlight, a broad H α line is obviously present (see Figure 1). To isolate the emission lines and measure their properties we modeled the continuum as a linear combination of starlight and a non-stellar component and subtracted it. To describe the starlight we experimented with spectra of elliptical or S0 galaxies (NGC 3379, NGC 4339, NGC 4365, and NGC 5322), while the non-stellar continuum was assumed to have a power-law spectrum of the form $f_\nu \propto \lambda^\alpha$. We found that the spectrum of NGC 4339 provides an excellent match to the continuum of NGC 3065, with no need for a non-stellar component. The non-stellar continuum need not contribute more than 10% of the flux at the wavelengths of H α or H β and no more than 15% of the flux at the wavelength of the [O II] $\lambda 3727$ line. The residual spectrum is shown in Figure 1 with the emission lines identified. It shows the broad H α line very clearly, as well as an unambiguously broad H β line. A hint of a broad H γ line is discernible as well. It is worth empha-

sizing that a careful subtraction of the starlight is needed to isolate the emission lines in the vicinity of H β since the starlight spectrum has a rather rich absorption-line structure in this region. It can easily hide or distort the appearance of the H β line, as well as the nearby [O III] $\lambda 4959$ line. Our approach to subtracting the starlight from the Mt. Hopkins spectrum was somewhat different, since the flux scale of this spectrum is not calibrated. We used as a template the spectrum of the S0 galaxy M42 (NGC 4472), observed during the CfA redshift with the same setup as and within 7 weeks of NGC 3065. We subtracted the template spectrum from that of NGC 3065, after normalizing the continuum around H α to unity in both spectra. This procedure resulted in the removal of all stellar absorption features from the spectrum of NGC 3065.

A lower limit to the equivalent width of the H α line, *relative to the non-stellar continuum* is $EW > 170$ Å. This limit is comparable to the equivalent widths of other LINERs with broad H α lines (e.g., NGC 4579, NGC 4450, NGC 4203, M81), which are in the range 140–530 Å. For comparison, the equivalent widths of the H α lines of well-known Seyfert galaxies (e.g., NGC 4151, NGC 5548, Mkn 6, Mkn 841; measured from the spectra of Eracleous & Halpern 1993) are somewhat larger, falling in the range of 500–600 Å.




FIG. 2.— Velocity-aligned profiles of the Balmer lines of NGC 3065 with the lines, observatories, and dates labelled. The vertical scale of the Mt. Hopkins spectrum is in normalized counts, while in all other cases it represents f_ν in mJy. The solid line in the top panel is a fit of a disk model to the broad line profile. The model is described in §4 of the text. In the third panel from the top we superpose the 1980 H α spectrum on the 2000 H α spectrum for comparison (the former spectrum has been scaled to match the [N II] $\lambda 6584$ strength of the latter; see §3 of the text for more details).

In Table 1 we list the relative emission-line intensities measured from the spectra after subtracting the continuum. They have been corrected for Galactic reddening using $E(B - V) = 0.067$ (Schlegel, Finkbeiner & Davis 1998). We also list the widths of the broad Balmer lines as well as the widths of the narrow lines included in the KPNO spectrum (unfortunately, the narrow lines in the MDM spectrum are not resolved). If we naively fit the broad Balmer lines with single Gaussian models, we find that their centroids are blueshifted by about 450 km s^{-1} from the reference frame defined by the narrow lines. However, we also note that at least the $H\alpha$ line shows a red asymmetry: the red wing extends about $2,000 \text{ km s}^{-1}$ further from the narrow component of the line than the blue wing does. To illustrate the properties of the broad-line profiles we plot $H\alpha$ and $H\beta$ spectra on a common velocity scale in Figure 2. This illustration brings out the features of the Balmer line profiles: a blueshifted shoulder and red wing that is more extended than the blue wing. It also shows that the broad $H\alpha$ line was absent in the 1980 spectrum, which means that it must have appeared some time in the past 20 years. To illustrate the difference between the 1980 and 2000 $H\alpha$ spectra more clearly, we superpose the two in the third panel of Figure 2. The 1980 spectrum has been scaled to match the $[\text{N II}] \lambda 6584$ strength of the 2000 spectrum. The difference between the two is easily discernible, especially on the blue side of the broad $H\alpha$ line, where the intensity in the 2000 spectrum is 4 times higher than what would be consistent with the noise of the 1980 spectrum.

TABLE 1
EMISSION-LINE PROPERTIES OF NGC 3065

Line	Intensity ^a	FWHM ^b (km s^{-1})
$[\text{O II}] \lambda 3727$	5.17	
$H\gamma$ (narrow)	0.58	
$H\gamma$ (broad)	3.25:	6,200
$H\beta$ (narrow)	1.00	
$H\beta$ (broad)	10.27	7,300
$[\text{O III}] \lambda 4959$	1.04	
$[\text{O III}] \lambda 5007$	3.09	
$[\text{O I}] \lambda 6300$	2.23	520
$[\text{O I}] \lambda 6363$	0.73:	
$[\text{N II}] \lambda 6548$	2.39	
$H\alpha$ (narrow)	2.72	375
$H\alpha$ (broad)	37.39	5,300
$[\text{N II}] \lambda 6583$	5.90	310
$[\text{S II}] \lambda 6717$	1.73	260
$[\text{S II}] \lambda 6731$	1.68	260

^a Line intensities are given relative to the *narrow* $H\beta$ line ($2.9 \times 10^{-15} \text{ erg cm}^{-2} \text{ s}^{-1} \pm 20\%$) after correction for Galactic reddening.

^b The line widths have been corrected for the instrumental resolution.

To assess whether the normalization procedure we adopted above is fair, we examined our 2-dimensional spectra of NGC 3065 to search for extended $[\text{N II}] \lambda 6584$ emis-

sion. If the $[\text{N II}] \lambda 6584$ flux were distributed along the slit, then the flux collected in the small apertures we used to extract the 2000 spectra would be considerably less than the flux collected in the larger aperture used in the 1980 observation. Our inspection showed the $[\text{N II}] \lambda 6584$ source to be unresolved. Thus the relative strengths of $[\text{N II}] \lambda 6584$ and broad $H\alpha$ are not affected by any missing $[\text{N II}] \lambda 6584$ flux and the above normalization procedure is valid.

4. RESULTS, DISCUSSION, AND SPECULATION

The relative intensities of its narrow lines make NGC 3065 a LINER. The oxygen line ratios, $[\text{O II}] \lambda 3727/[\text{O III}] \lambda 5007 = 1.6$ and $[\text{O I}] \lambda 6300/[\text{O III}] \lambda 5007 = 0.7$, satisfy Heckman's original definition of the class. Other line ratios, such as $[\text{O III}] \lambda 5007/H\beta = 3.3$, $[\text{N II}] \lambda 6583/H\alpha = 2.2$, $[\text{O I}] \lambda 6300/H\alpha = 0.8$, and $[\text{S II}] \lambda \lambda 6717, 6731/H\alpha = 1.3$, place NGC 3065 in the regions of the diagnostic ratio diagrams occupied by LINERs, albeit close to the boundary with Seyferts (see, for example, Ho et al. 1997a). The widths of the narrow forbidden lines appear to follow a trend with critical density: lines of higher critical density are broader than lines of lower critical density. Such a trend is often observed in LINERs and it has been interpreted as an indication that the narrow-line emitting gas is stratified in density (Filippenko 1985). The presence of broad Balmer lines in the spectrum of NGC 3065 establishes it as an AGN beyond doubt, and confirms the suggestion of Iyomoto et al. (1998) based on the X-ray properties.

Finding broad Balmer lines in the spectra of LINERs is not unusual. For example, M81 has been known for quite some time to have broad Balmer lines (Peimbert & Torres-Peimbert 1981; Filippenko & Sargent 1988). More recently, broad Balmer lines have been found in several LINERs with the *Hubble Space Telescope* (*HST*) which can obtain spectra through very small apertures that exclude the contaminating starlight very effectively. Examples include NGC 4203 (Shields et al. 2000), NGC 4450 (Ho et al. 2000), and NGC 4579 (Barth et al. 2000). The sudden appearance or dramatic variability of broad Balmer lines in LINERs is not unheard of either: it has been observed in at least two other cases so far, NGC 1097 (Storchi-Bergmann et al. 1993) and M81 (Bower et al. 1996). It is also possible that the broad $H\alpha$ lines of NGC 4203, NGC 4450, and NGC 4579 varied dramatically between the early observations from the mid-1980s (Ho et al. 1997b) and the later *HST* observations from the late 1990s, although one cannot be confident in view of the available data.

The broad Balmer lines of LINERs often have double-peaked profiles, which are characteristic of rotation and suggest an origin in an accretion disk around a supermassive black hole (see above references). This underscores an intimate connection between LINERs and another class of double-peaked emission line AGNs, the broad-line radio galaxies (hereafter BLRGs; Eracleous & Halpern 1994), which is bolstered by other similarities between the two classes of object. In particular, the relative strengths of the narrow emission lines of BLRGs with double-peaked emission lines approach those of LINERs (the prototype, Arp 102B is a certified LINER; Stauffer, Schild, & Keel 1983). Moreover, in Pictor A, whose relative narrow-line

strengths are close to LINER-like (Filippenko 1985), the double-peaked Balmer lines appeared abruptly in the mid-to late-1980s (Halpern & Eracleous 1994; Sulentic et al. 1995).

Although the broad Balmer lines of NGC 3065 are not double peaked, the fact that their red wing extends further than the blue wing is reminiscent of gravitational redshift of photons originating in the inner part of an accretion disk. We thus speculate that the broad Balmer lines of NGC 3065 originate in the outer parts of an accretion disk around a supermassive black hole. We emphasize that this is by no means a unique explanation for the origin of the broad Balmer lines; it is merely inspired by the profiles of the Balmer lines of other similar objects. To assess the plausibility of this hypothesis we have tried to fit their profiles with the disk model developed by Chen, Halpern, & Filippenko (1989) and Chen & Halpern (1989). The result of this exercise is superposed on the observed H α profile shown in the top panel of Figure 2. According to the adopted model, the axis of the disk is inclined at an angle of 50° to the line of sight and the line-emitting region is between radii of 900 and 100,000 GM/c^2 , where M is the mass of the black hole. The emissivity is a broken power law with radius: $\epsilon \propto r^{-q}$, with $q = 1.7$ for $r < 10,000 GM/c^2$ and $q = 3$ elsewhere. In this context it is noteworthy that the profiles of emission lines coming from an accretion disk need not be double-peaked: if the ratio of the inner-to-outer radius of the line-emitting part of the disk is large or the disk is close to face on, the two peaks get close enough together that they merge and the profile appears flat-topped or single-peaked (Eracleous 1999; Corbin 1997). Other combinations of model parameters may be able to produce equally good fits. We have not explored the parameter space because we are only focusing on the plausibility of this interpretation here. Also, more sophisticated disk models which include an eccentricity (Eracleous et al. 1995) or a spiral wave (Gilbert et al. 1999) may be able to reproduce the observed H α profile even better, but such detailed modeling is well outside the scope of this paper.

To further explore the similarity of NGC 3065 with LINERs and BLRGs with double-peaked Balmer lines we have estimated the mass of its central black hole and the corresponding Eddington luminosity. To estimate the black hole mass we used the recently established correlation between it and the stellar velocity dispersion in the host galaxy (Ferrarese & Merrit 2000; Gebhardt et al. 2000). The stellar velocity dispersion of $173 \pm 16 \text{ km s}^{-1}$ reported by Tonry & Davis (1981) thus yielded $M_\bullet = (9 \pm 4) \times 10^7 M_\odot$, where the error bar reflects not only the uncertainty in the velocity dispersion but also uncertainties in the parameters describing the correlation between the velocity dispersion and the black hole mass (see, Ferrarese & Merrit 2000; Gebhardt et al. 2000). As a check, we also estimated the black hole mass based on the correlation between it and the blue luminosity of the bulge of the host galaxy (see the latest version in Kormendy 2000) and using the bulge-disk decomposition of Kormendy (1977), obtaining $M_\bullet = 2 \times 10^8 M_\odot$. This value is almost a factor of 2 higher than that obtained with the previous method, which is very likely a result of the large scatter about the mean trend between the

black hole mass and the bulge luminosity. Because of this large dispersion, we prefer the black hole mass inferred from the stellar velocity dispersion. The implied Eddington luminosity is $L_{\text{Edd}} = 1.3 \times 10^{46} \text{ erg s}^{-1}$. If we take the bolometric accretion luminosity of NGC 3065 to be 10 times larger than the observed 2–10 keV X-ray luminosity (see, for example, Ho 1999), we find an Eddington ratio of $L_{\text{bol}}/L_{\text{Edd}} \approx 2 \times 10^{-4}$, which indicates a very low relative accretion rate, a common feature of LINERs (see Ho 1999). At such a low accretion rate the inner accretion disk is likely to be “advection dominated” (an ADAF or ion torus; Narayan & Yi 1994, 1995; Rees et al. 1982) and to form a vertically extended structure that can illuminate the outer, thin disk effectively. Thus, it could power the observed broad-line emission (cf, Chen & Halpern 1989). This geometrical requirement may very well be the reason why disk-like emission lines are preferentially found in AGNs with very low accretion rates relative to the Eddington rate. In fact, if we are to associate the broad Balmer lines of NGC 3065 with emission from an accretion disk, then external illumination of the disk is needed in order to power the line emission. This is because the H α luminosity is $3 \times 10^{40} \text{ erg s}^{-1}$ while the viscous power output of the line-emitting portion of the disk (calculated following Eracleous & Halpern 1994) is only $4 \times 10^{39} \text{ erg s}^{-1}$. Yet another appealing feature of ADAFs in LINERs is the fact that their hard spectral energy distribution, which lacks a “UV bump,” when combined with a low ionization parameter, can explain the relative strengths of the narrow emission lines (Halpern & Steiner 1983; Ferland & Netzer 1983).

The reason for the recent emergence of the broad Balmer lines in NGC 3065, as well as in similarly behaved objects, remains an open question. One possibility is that the emission lines come from a transient accretion disk which formed from the debris released by the tidal disruption of a star by the black hole (cf, NGC 1097; Eracleous et al. 1995; Storchi-Bergmann et al. 1995). Another is a change in the structure of the inner accretion disk associated with a change in the accretion rate, i.e., a transformation from a thin disk to an ADAF (Storchi-Bergmann et al. 1997). Perhaps the long-term variations of the Balmer-line profiles will provide clues to their origin. The variations may show evidence for dynamical phenomena (e.g., spiral waves in the disk; Gilbert et al. 1999), which may cause fluctuations in the accretion rate. Alternatively, the variations may show evidence for a disk geometry that can be related to its formation process (e.g., an eccentric disk formed from tidal debris; Eracleous et al. 1995). We will continue to monitor the broad Balmer lines of NGC 3065 in an effort to uncover their cause.

We thank J. Huchra for sending us the spectra of NGC 3065 and NGC 4472 obtained during the CfA Redshift Survey. We are grateful to Luis Ho, Aaron Barth, and the anonymous referee for their insightful comments and suggestions. In our investigation we have made use of the NASA/IPAC Extragalactic Database (NED) which is operated by the Jet Propulsion Laboratory, California Institute of Technology, under contract with the National Aeronautics and Space Administration.

REFERENCES

- Barth, A. J., Ho, L. C., Filippenko, A. V., Rix, H.-W., & Sargent, W. L. W. 2000, *ApJ*, in press (astro-ph/0008273)
- Bower, G. A., Wilson, A. S., Heckman, T. M. & Richstone, D. O. 1996 *AJ*, 111, 1901
- Burbidge, E. M. & Burbidge, G. R. 1965, *ApJ*, 142, 634
- Chen, K. & Halpern, J. P. 1989, *ApJ*, 344, 115
- Chen, K., Halpern, J. P., & Filippenko, A. V. 1989, *ApJ*, 339, 742
- Condon, J. J., Cotton, W. D., Greisen, E. W., Yin, Q. F., Perley, R. A., & Taylor, G. B. 1998, *ApJ*, 115, 1693
- Corbin, M. R. 1997, *ApJ*, 485, 517
- Eracleous, M. 1999, in "Structure and Kinematics of Quasar Broad Line Regions," ASP Conference Series, Vol. 175. eds. C. M. Gaskell, et al. (San Francisco: ASP), 163
- Eracleous, M. & Halpern, J. P. 1993, *ApJ*, 409, 584
- Eracleous, M. & Halpern, J. P. 1994, *ApJS*, 90, 1
- Eracleous, M., Livio, M., Halpern, J. P., & Storchi-Bergmann, T. 1995, *ApJ*, 438, 610
- Fabbiano, G., Kim, D.-W., & Trinchieri, G. 1992, *ApJS*, 80, 531
- Ferland, G. J. & Netzer, H. 1983, *ApJ*, 264, 105
- Ferrarese, L. & Merrit, D. 2000, *ApJ*, 539, L9
- Filippenko, A. V. 1996, "The Physics of LINERs in View of Recent Observations," ASP Conference Series, Vol. 103; eds. M. Eracleous et al. (San Francisco: ASP), 17
- Filippenko, A. V. 1985, *ApJ*, 289, 475
- Filippenko, A. V. & Sargent, W. L. W. 1988, *ApJ*, 324, 134
- Gebhardt, K., et al. 2000, *ApJ*, 539, L13
- Gilbert, A. M., Eracleous, M., Filippenko, A. V., & Halpern J. P. 2000, in "Structure and Kinematics of Quasar Broad Line Regions, ASP Conference Series," Vol. 175. eds. C. M. Gaskell, et al. (San Francisco: ASP), 189
- Halpern, J. P. & Eracleous, M. 1994, *ApJ*, 433, L17
- Halpern, J. P. & Steiner, J. E. 1983, *ApJ*, 269, L37
- Heckman, T. M. 1980, *A&A*, 87, 152
- Ho, L. C. 1999, *ApJ*, 516, 672
- Ho, L. C., Filippenko, A. V., & Sargent, W. L. W. 1997a, *ApJS*, 112, 315
- Ho, L. C., Filippenko, A. V., & Sargent, W. L. W., & Peng, C. Y. 1997b, *ApJS*, 112, 391
- Ho, L. C., Rix, H.-W., Shields, J. C., Rudnick, G., McIntosh, D. H., Filippenko, A. V., & Sargent, W. L. W., & Eracleous, M. 2000, *ApJ*, 541, 120
- Humason, M. L., Mayall, N. U., & Sandage, A. R. 1956, *AJ*, 61, 97
- Iyomoto, N., Makishima, K., Matsushita, K., Fukazawa, Y., Tashiro, M. & Ohashi, T. 1998, *ApJ*, 503, 168
- Kormendy, J. 1977, *ApJ*, 217, 406
- Kormendy, J. 2000, in "Galaxy Disks and Disk Galaxies," eds. J. G. Funes & E. M. Corsini (San Francisco: ASP), in press (astro-ph/0007401).
- Narayan, R. & Yi, I. 1994, *ApJ*, 443, L41
- Narayan, R. & Yi, I. 1995, *ApJ*, 444, 231
- Peimbert, M. & Torres-Peimbert, S. 1981, *ApJ*, 245, 845.
- Ptak, A., Serlemitsos, P., Yaqoob, T., & Mushotzky, R. F. 1999, *ApJS*, 120, 179
- Rees, M. J., Begelman, M. C., Blandford, R. D., & Phinney, E. S. 1982, *Nature*, 295, 17
- Schlegel, D., Finkbeiner, D. P., & Davies, M. 1998, *ApJ*, 500, 525
- Shields, J. C., Rudnick, G., Rix, H.-W., Ho, L. C., McIntosh, D. H., Filippenko, A. V., & Sargent, W. L. W. 2000, *ApJ*, 534, L27
- Storchi-Bergmann, T., Baldwin, J. A., & Wilson, A. S. 1993, *ApJ*, 410, L11
- Storchi-Bergmann, T., Eracleous, M., Livio, M., Wilson, A. S., Filippenko, A. V., Halpern, J. P. 1995, *ApJ*, 443, 617
- Storchi-Bergmann, T., Eracleous, M., Ruiz, M. T., Livio, M., Wilson, A. S., & Filippenko, A. V. 1997, *ApJ*, 489, 87
- Stauffer, J., Schild, R., & Keel, W. C. 1983, *ApJ*, 270, 465
- Sulentic, J. W., Marziani, P., Zwitter, T., Calvani, M. 1995, *ApJ*, 438, L1
- Terashima, Y., Ho, L. C., Ptak, A. F., Mushotzky, R. F., Serlemitsos, P., Yaqoob, T., & Kunieda, H. 2000, *ApJ*, 533, 729
- Tonry, J. L. & Davis, M. 1981, *ApJ*, 246, 666
- Tully, R. B. 1987, *Nearby Galaxies Catalog* (Cambridge: Cambridge University Press)